

ACL Reconstruction with Cadaver tissue

Hussein Elkousy, MD
Fondren Orthopedic Group
7401 South Main St.
(713) 799-2300

The problem:

Tear of the anterior cruciate ligament (ACL).

Reason for treatment:

The knee buckles or gives way with daily activities, work, or recreational activities. The ligament tear usually occurs from a specific injury during work, a fall, or a sporting activity.

The treatment:

The torn ligament is replaced with a tendon from another person who has donated their tissue. The new ligament is held in place using pins and screws.

Day of surgery:

The patient should plan to arrive 2 hours prior to the procedure to get situated and meet the operative team. The procedure is done arthroscopically, but a small open incision is required to place the bone tunnels. This incision is between 1 and 2 inches long and is on the front and inside of the knee. Tunnels are drilled through the tibia and femur bones to place the new ligament. The pins and screws are placed inside the bone.

The patient is asleep for the procedure. The procedure itself takes between 60 and 90 minutes depending on whether any other surgery is necessary on the cartilage. Recovery room time is approximately 1 hour. Patients are encouraged to stay in the hospital overnight for pain control but many go home the same day.

After surgery:

The patient is generally allowed to bear partial weight with a brace on and locked out with the knee straight. Crutches are used for balance and support. The brace may only be removed when bathing or when the patient is awake and sitting/lying. It may not be removed if the patient is moving from one area to another.

Exercises are begun the day of surgery to bend the knee and strengthen the knee with the brace off, but not while the patient is standing. These exercises should be done daily until clinic follow-up in 1-2 weeks. A machine is used which bends the knee for you initially. This is called a CPM machine (Continuous Passive Motion).

The knee is often painful, swollen, and warm for several weeks after surgery. This can be managed with icing the knee several times each day (20 minutes each time), using a compressive wrap (ACE), and taking pain medication provided by a prescription. Do not take any anti-inflammatory medications (Advil, Alleve, etc.) for 6 weeks after the surgery. Most patients will be able to stop using pain medicine within 2 weeks.

By six weeks, most patients walk normally without the brace. The knee is still protected for 3-6 months. Most patients can return to playing sports in 6 to 8 months.